Pharmasin®

100% W/W tylosin tartrate
Water soluble granules





Pure and powerful

HUVEPHARMA®
We add performance to your business

PHARMASIN® 100% W/W WATER SOLUBLE GRANULES

Composition

1.1 g of granules contains 1 g of tylosin (corresponding to 1.1 g of tylosin tartrate)

Pharmaceutical form

Granules for use in drinking water.

Target species

Calves, pigs, chickens, turkeys

Indications

Calves

- Treatment and prevention of pneumonia caused by *Mycoplasma spp* when the disease has been established in the herd.

Pigs

- Treatment and prevention of enzootic pneumonia caused by *Mycoplasma hyopneumoniae* and *Mycoplasma hyorhinis* when the disease has been established in the herd.
- Treatment and prevention of *Porcine Intestinal Adenomatosis* (Ileitis) associated with *Lawsonia intracellularis* when the disease has been established in the herd.

Chickens

- Treatment and prevention of chronic respiratory diseases (CRD) caused by *Mycoplasma gallisepticum* and *Mycoplasma synoviae* when the disease has been established in the flock.
- Treatment and prevention of necrotic enteritis caused by *Clostridium perfringens* when the disease has been established in the flock.

Turkeys

- Treatment and prevention of infectious sinusitis caused by *Mycoplasma gallisepticum* when the disease has been established in the flock.

Practical dosing and administration

Oral administration through the drinking water. In calves the product can also be administered through milk or milk replacer.

Calves

- 10-20 mg tylosin per kg BW (corresponding to 11-22 mg of the veterinary medicinal product per kg BW), twice daily (corresponding to a daily dose of 20-40 mg tylosin per kg BW), for 7-14 days.

Pigs

- For the treatment of enzootic pneumonia: 20 mg tylosin per kg BW per day (corresponding to 22 mg of the veterinary medicinal product per kg BW) for 10 days.
- For the treatment of ileitis or PIA: 5-10 mg tylosin per kg BW per day (corresponding to 5.5-11 mg of the veterinary medicinal product per kg BW) for 7 days.

Chickens

- For the treatment of chronic respiratory disease: 75-100 mg tylosin per kg BW per day (corresponding to 82.5-110 mg of the veterinary medicinal product per kg BW) for 3-5 days.
- For the treatment of necrotic enteritis: 20 mg tylosin per kg BW per day (corresponding to 22 mg of the veterinary medicinal product) for 3 days.

Turkeys

- 75-100 mg tylosin per kg BW per day (corresponding to 82.5-110 mg of the veterinary medicinal product per kg BW) for 3-5 days.

For the preparation of the medicated water/milk/milk-replacer the body weight of the animals to be treated and their actual daily water/milk/milk-replacer consumption should be taken into due account. Consumption may vary depending on factors like age, state of health, breed, husbandry system. To provide the required amount of active substance in mg per litre drinking water/milk/milk-replacer the following calculation should be made:

$$\frac{\text{...mg tylosin per kg bodyweight per day} \times \text{Average bodyweight (kg) of the animals to be treated}}{\text{Average amount of drinking water or milk/animal (l)}} = \text{...mg tylosin/l of drinking water}$$

Withdrawal period(s)

- Calves (meat and offal): 12 days
- Pigs (meat and offal): 1 day
- Turkeys (meat and offal): 2 days
- Turkey (eggs): 0 days
- Chickens (meat and offal): 1 day
- Chicken (eggs): 0 days

Shelf-life

- Shelf-life of the veterinary medicinal product as packaged for sale: 3 years
- Shelf-life after dilution or reconstitution according to directions:
  - medicated water: 24 hours
  - medicated milk or milk replacer: 24 hours
- Shelf-life after first opening the immediate packaging: 3 months

Packaging

- 1.1 kg resealable block bottom zipped sachet made of polyethylene/aluminium/polyethylene terephthalate laminate
- 110 g high density polyethylene pot with polypropylene cap



Contraindications

Do not use in animals with known hypersensitivity to tylosin or other macrolides.
Do not use in cases of known resistance to tylosin or cross-resistance to other macrolides (MLS-resistance).
Do not use in animals vaccinated with tylosin-sensitive vaccines either at the same time or within 1 week previously.
Do not use in animals with hepatic disorders.
Do not use in horses.

- Full SPC can be obtained on request.
- Indications listed above are not necessarily authorized in all countries. Please consult the local label for exact indications and posology.
- Use medicines responsibly.
- POM-V
- Vm number 30282/4017

HUVEPHARMA®

Pharmasin®

Phar100.WSG. Calf_Turkey_Pigs&Poultry. short.Br. EN02. 0918/Novis

HUVEPHARMA EOOD • 3A NIKOLAY HAYTOV STR • 1113 SOFIA • **BULGARIA** • P +359 2 862 5331 • F +359 2 862 5334 • SALES@HUVEPHARMA.COM
HUVEPHARMA N.V. • UITBREIDINGSTRAAT 80 • 2600 ANTWERP • **BELGIUM** • P +32 3 288 1849 • F +32 3 289 7845 • CUSTOMERSERVICE@HUVEPHARMA.COM