
Tilmovet®

250 mg tilmicosin/ ml
Concentrate for oral solution



Best choice for
respiratory infections

HUVEPHARMA®
We add performance to your business

TILMOVET® 250 MG/ ML CONCENTRATE FOR ORAL SOLUTION

Composition
Active substance: **250 mg tilmicosin per ml**

Pharmaceutical form
Concentrate for oral solution

Target species
Pigs, chickens (broilers and pullets), turkeys and cattle (calves)

Indications
Pigs
- Treatment and prevention of respiratory infections associated with *Mycoplasma hyopneumoniae, Pasteurella multocida* and *Actinobacillus pleuropneumoniae* when the disease has been diagnosed at the herd level.

Chickens
- Treatment and prevention of respiratory infections in poultry flocks associated with *Mycoplasma gallisepticum* and *Mycoplasma synoviae* when the disease has been diagnosed at the herd level.

Turkeys
- Treatment and prevention of respiratory infections in turkey flocks associated with *Mycoplasma gallisepticum* and *Mycoplasma synoviae* when the disease has been diagnosed at the herd level.

Calves
- Treatment and prevention of respiratory infections associated with *Mannheimia haemolytica, Pasteurella multocida, Mycoplasma bovis* and *Mycoplasma dispar* when the disease has been diagnosed at the herd level.

Practical dosing and administration
- For oral use only.
- The product must be diluted in drinking water or milk replacer before administration.

Pigs
- 15-20 mg tilmicosin per kg BW
- = 6-8 ml product for 100 kg BW
- = 80 ml product per 100 l drinking water
- For 5 days

Chickens
- 15-20 mg tilmicosin per kg BW
- = 6-8 ml product for 100 kg BW
- = 30 ml product per 100 l drinking water
- For 3 days

Turkeys
- 10-27 mg tilmicosin per kg BW
- = 4-11 ml product for 100 kg BW
- = 30 ml product per 100 l drinking water

Calves
- 12.5 mg tilmicosin per kg BW 2 times per day
- = 1 ml product for 20 kg BW 2 times per day
- For 3-5 days
  One 960 ml bottle is sufficient to medicate
- 1200 l drinking water for pigs
- 3200 l drinking water for broilers, turkeys and pullets
- Drinking water or milk replacer for 48-80 calves (40 kg BW)

Withdrawal period(s)
Meat and offal
- Pigs: 14 days
- Chickens: 12 days
- Turkeys: 19 days
- Calves: 42 days
  Eggs: not authorised for use in birds producing eggs for human consumption

Shelf life
- Shelf life of the veterinary medicinal product as packaged for sale: 2 years
- Shelf life after first opening the immediate packaging: 3 months
- Shelf life after dilution in drinking water: 24 hours
- Shelf ife after reconstitution in milk replacer: 4 hours

Packaging
960 ml is presented in a white high density polyethylene bottle with white polypropylene or high density polyethylene, tamper-evident cap.
HUVEPHARMA



Contraindications
Do not use in case of hypersensitivity to the active substance or in cases of known resistance to tilmicosin. Do not use in horses.

- Full SPC can be obtained on request.
- Indications listed above are not necessarily authorized in all countries. Please consult the local label for exact indications and posology.
- Use medicines responsibly.
- POM-V
- Vm number 30282/4001

Tilmovet®

Tilmovet 250.OS.AllSp.Br.EN01.1218./Novis

HUVEPHARMA®

HUVEPHARMA EOOD • 3a NIKOLAY HAYTOV STR. • 1113 SOFIA • **BULGARIA** • P +359 2 862 5331 • F +359 2 862 5334 • SALES@HUVEPHARMA.COM
HUVEPHARMA N.V. • UITBREIDINGSTRAAT 80 • 2600 ANTWERP • **BELGIUM** • P +32 3 288 1849 • F +32 3 289 7845 • CUSTOMERSERVICE@HUVEPHARMA.COM